


# The Niels Henrik Abel Contest 1993

## FINAL

### Problem 1

a) Let  $ABCD$  be a convex quadrilateral. (I.e. the angles are all less than  $180^\circ$ .) Let  $A'$  be the midpoint of  $AB$ ,  $B'$  the midpoint of  $BC$ ,  $C'$  the midpoint of  $CD$ , and  $D'$  the midpoint of  $AD$ . Draw the lines  $A'C'$  and  $B'D'$ , and let  $a, b, c$ , and  $d$  be the areas of the four minor quadrilaterals, as shown in the figure. Prove that  $a + c = b + d$ .



b) Given a triangle with sides of length  $a, b$ , and  $c$ , prove that

$$\frac{a}{b+c} + \frac{b}{a+c} + \frac{c}{a+b} < 2.$$

### Problem 2

Prove that if  $b < c < d$ , then the inequality

$$(a + b + c + d)^2 > 8(ac + bd)$$

holds for all  $a$ .

### Problem 3

The Fermat-numbers are defined by  $F_n = 2^{2^n} + 1$  for  $n = 0, 1, 2, \dots$

a) Prove that  $F_n = F_{n-1}F_{n-2} \cdots F_1F_0 + 2$  for  $n = 1, 2, 3, \dots$

b) Prove that two different Fermat-numbers cannot have a common factor greater than 1.

### Problem 4

We have a cube. Each of the 8 corners are given values 1 or  $-1$ . Each of the six sides are given a value equaling the product of its four corners. Let  $A$  be the sum of the values of all the corners (eight) and all the sides (six). Then,  $A$  is the sum of 14 values. Which values of  $A$  are attainable?