

The 37th International Mathematical Olympiad

First day (10 July, 1996)

Time: $4\frac{1}{2}$ hours

Version: English

1. Let $ABCD$ be a rectangular board with $|AB| = 20$, $|BC| = 12$. The board is divided into 20×12 unit squares. Let r be a given positive integer. A coin can be moved from one square to another if and only if the distance between the centres of the two squares is \sqrt{r} . The task is to find a sequence of moves taking the coin from the square which has A as a vertex to the square which has B as a vertex.
 - (a) Show that the task cannot be done if r is divisible by 2 or 3.
 - (b) Prove that the task can be done if $r = 73$.
 - (c) Can the task be done when $r = 97$?
2. Let P be a point inside triangle ABC such that

$$\angle APB - \angle ACB = \angle APC - \angle ABC.$$

Let D , E be the incentres of triangles APB , APC respectively. Show that AP , BD and CE meet at a point.

3. Let $S = \{0, 1, 2, 3, \dots\}$ be the set of non-negative integers. Find all functions f defined on S and taking their values in S such that

$$f(m + f(n)) = f(f(m)) + f(n) \text{ for all } m, n \text{ in } S.$$

Each problem is worth 7 points.

The 37th International Mathematical Olympiad

Second day (11 July, 1996)

Time: $4\frac{1}{2}$ hours

Version: English

4. The positive integers a and b are such that the numbers $15a + 16b$ and $16a - 15b$ are both squares of positive integers. Find the least possible value that can be taken by the minimum of these two squares.
5. Let $ABCDEF$ be a convex hexagon such that AB is parallel to ED , BC is parallel to FE and CD is parallel to AF . Let R_A, R_C, R_E denote the circumradii of triangles FAB, BCD, DEF respectively, and let p denote the perimeter of the hexagon. Prove that

$$R_A + R_C + R_E \geq \frac{p}{2}.$$

6. Let n, p, q be positive integers with $n > p + q$. Let x_0, x_1, \dots, x_n be integers satisfying the following conditions:
- (a) $x_0 = x_n = 0$;
- (b) for each integer i with $1 \leq i \leq n$,

$$\text{either } x_i - x_{i-1} = p \text{ or } x_i - x_{i-1} = -q.$$

Show that there exists a pair (i, j) of indices with $i < j$ and $(i, j) \neq (0, n)$ such that $x_i = x_j$.

Each problem is worth 7 points.